نموذج استرشادى (٥) لامتحان شهادة إتمام الدراسة الثانوية العامة ٢٠٢٥ / ٢٠٢٦ م

الرياضيات التطبيقية (الشعبة العلمية – رياضيات) الزمن : ساعتان

First: Multiple choice questions" one mark for each item"

(1) If the force $\overrightarrow{F} = 3\vec{\imath} + 4\vec{\jmath}$ acts at the point (2,3) then the length of the perpendicular drawn from the point (-1,3) to the line of action of the force = ……. length units

(a) 1.2 (b) 1.8 (c) 2.4 (d) 4.8

(2) In the opposite figure : forces of magnitudes F , 5 , k and 7 Newtons act at the points A,C,D and B respectively , if the forces are parallel and in equilibrium then K-F=…….



(a) 1 (b) 2 (c) 3 (d) 4

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(3) In the opposite figure: ABCD is a square of side length 8cm, if the two forces (F, F) form couple equivalent to the couple formed by the two forces (20,20) where the two forces $\overrightarrow{F}$, $\overrightarrow{F}$ are parallel to $\overleftrightarrow{BD}$, then F =…… Newton



(a) $5\sqrt{2}$ (b) $10\sqrt{2}$ (c) $15\sqrt{2}$ (d) $20\sqrt{2}$

(4) If the velocity vector of a particle $\overrightarrow{v}$(m/s) at any time t(sec) is given by the relation $\overrightarrow{v} = (t^2 - 6t + 5)\overrightarrow{c}$, then the magnitude of the acceleration after 5 seconds from starting the motion =…… m/s^2

(a) 3 (b) 4 (c) 5 (d) 6

(5) If the mass of a body m (kg) at any time t (sec) is given by the relation m =(2t+5) kg moves with uniform velocity 49 m/s, then the magnitude of the force acting on it =……N.

(a) 28 (b) 49 (c)70 (d) 98

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(6) In the opposite figure: ABCD is a rectangle in which AB=3cm , if the forces are in equilibrium, then AD=…….cm



(a) 3 (b) 4 (c) 5 (d) 6

(7) In the opposite figure: ABCD is a rhombus of side length 4cm where BD=AB , if the system of the forces(measured in Newton) form couple then the magnitude of its moment=…… Newton.cm



(a) $10\sqrt{3}$ (b) $15\sqrt{3}$ (c) $20\sqrt{3}$ (d) $25\sqrt{3}$

(8) A particle moves in a straight line where its velocity vector $\overrightarrow{v}$(m/s) at any time t(sec) is given by the relation $\overrightarrow{v} = (3t^2 + 2t)\overrightarrow{c}$, then the algebraic measure of its displacement during the time interval $[3,6]$ = …… meters

(a) -216 (b) -87 (c) 87 (d) 216

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(9) If a man of weight 70 kg.wt stands on the floor of a lift moving vertically up with acceleration 1.4 m/s^2 , then the pressure of the man on the floor of the lift =…….kg.wt

(a) 70 (b) 80 (c) 686 (d) 784

(10) A body moves in a straight line under the effect of a force $\overrightarrow{F}$ where its magnitude at any position "x" by the relation $F= 3x^2 - 2x$. If the force acts in the same direction of motion, then the work done by the force from the position x = 2 to the position x = 5 equals …….. work units

(a) 57 (b) 73 (c) 96 (d) 104

Second : Multiple choice questions" two marks for each item"

(11) The opposite figure : $\overline{AB}$ is a rod of negligible weight, its length 100cm, the drawn forces (measured by the same unit) act in the rod . If the rod is in equilibrium under the effect of these forces , then CD=……cm



(a) $10\sqrt{3}$ (b) $15\sqrt{3}$ (c) $20\sqrt{3}$ (d) $25\sqrt{3}$

(12) If the force $\overrightarrow{F} = 3\vec{\imath} - 4\vec{\jmath}$ (Newton) acted on a body so it moved from the point A(1,3) to the point B(7,3) , then the change in the potential energy of the body = …… work units

(a) -24 (b) -18 (c) 18 (d) 24

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(13) ABCD is a rectangle, the line of action of the force $\overrightarrow{F}$ passing through the center of the rectangle, if the moment vector of the force $\overrightarrow{F}$ about the point A = $-6\vec{k}$, then the moment vector of the force about the point C =…….$\vec{k}$

(a) -6 (b) -3 (c) 3 (d) 6

(14) If the forces $\overrightarrow{F_1} = 3\vec{\imath} + 4\vec{\jmath}$, $\overrightarrow{F_2} = \vec{\imath} - 2\vec{\jmath}$, $\overrightarrow{F_3} = -3\vec{\imath} + 4\vec{\jmath}$ act at the point A(2,3) then the moment vector of the resultant of the forces about the point B(1,1)=…….. $\vec{k}$

(a) -13 (b) -4 (c) 4 (d) 13

(15) If the opposite figure represents (velocity-time) graph for a body moves in a straight line, then its displacement after 8 seconds from starting the motion =……. cm



(a) -120 (b) 0 (c) 60 (d) 120

(16) A body of mass 10 kg rests on a horizontal smooth plane, if a horizontal force of magnitude 40 Newton acted on it , the body moved in its direction, then the magnitude of the velocity of the body after 5 seconds from starting its motion = ……m/s

(a) 10 (b) 20 (c) 30 (d) 40

(17) In the opposite figure: a uniform ladder its weight (w) rests with its upper end on a smooth vertical wall and its lower end on a rough horizontal ground , if the ladder is about to slide when it inclined angle of measure 45^0 to horizontal , then the coefficient of static friction between the ladder and the ground =……

حقوق النشر محفوظة لمديرية التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(a) $\frac{\sqrt{2}}{2}$ (b) $\frac{1}{2}$ (c) $\frac{2}{3}$ (d) $\sqrt{2}$

(18) The opposite figure represents the (force- displacement) graph for a body moving in a straight line ,if the force measured in Newton and the displacement in meter , then the change in kinetic energy of the body when the magnitude of the displacement =10 meters equals.....joule



(a) 110 (b) 115 (c) 120 (d) 125

Third: essay questions “two marks for each question”.

(19) In the opposite figure: ABC is right angled triangle at B in which AB=3cm , BC=8cm , force F(Newton) acts in the plane of the triangle such that $\overrightarrow{M_A} = \vec{0}$, $\overrightarrow{M_C} = -\overrightarrow{M_B} = 60\vec{k}$, Find $\left\|\overrightarrow{F}\right\|$.

A

3cm

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(20) A body is placed at the top of an inclined smooth plane of height 10 meter, find the velocity of the body when it reached the bottom of the plane

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

موقع الدكتور محمد رزق التعليمي

امسح الباركود بالموبيل لتثبيت التطبيق وتحميل كل الملازم مجانا وبسرعه

MR

حمل التطبيق الأن

GET IT ON Google Play

ملازم ابتدائي | ملازم اعدادي | ملازم ثانـــوي | كورسات ونتائج امتحانات

موقع الدكتور محمد رزق
Dr.MoRizK

اجابه الامتحان الاسترشادي الخامس رياضه تطبيقيه لغات انجليزي 2026 موقع الدكتور محمد رزق التعليمي

رقم السؤال	الاجـــــــابة
1	C
2	B
3	B
4	B
5	D
6	B

تم بواسطه موقع الدكتور محمد رزق التعليمي

7	C
8	D

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK

9	B
10	C
11	D
12	B
13	D
14	C
تم بواسطه موقع الدكتور محمد رزق التعليمي	
15	B
16	B
17	B
18	B

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK

19 In the opposite figure: ABC is right angled triangle at B in which AB=3cm , BC=8cm , force F(Newton) acts in the plane of the triangle such that $\overrightarrow{M_A} = \vec{O}$, $\overrightarrow{M_C} = -\overrightarrow{M_B} = 60\,\vec{k}$ Find $\| \vec{F} \|$. $\overrightarrow{M_A} = \vec{O}$ $\therefore$ A lies on line of action of $\vec{F}$ $\overrightarrow{M_C} = -\overrightarrow{M_B}$ line of action of $\vec{F}$ bisects $\overline{BC}$ 	19
19 In the opposite figure: ABC is right angled triangle at B in which AB=3cm , BC=8cm , force F(Newton) acts in the plane of the triangle such that $\overrightarrow{M_A} = \vec{O}$, $\overrightarrow{M_C} = -\overrightarrow{M_B} = 60\,\vec{k}$ Find $\| \vec{F} \|$. $\overrightarrow{M_B} = -\overrightarrow{60}$ $-F \times \frac{3 \times 4}{5} = -60$ $F = 25\ N$ 	
20 A body is placed at the top of an inclined smooth plane of height 10 meter, find the velocity of the body when it reached the bottom of the plane $T_0 + P_0 = T_1 + P_1$ $0 + m \times 9.8 \times 10 = 0 + \frac{1}{2} \times m \times v^2$ $9.8 \times 10 \times 2 = v^2$ $196 = v^2$ $v = 14m/sec$ 	20

تم الاجابه من خلال موقع الدكتور محمد رزق التعليمي

والمصدر قناه مدرستنا 3

موقع الدكتور محمد رزق التعليمي

امسح الباركود بالموبيل لتثبيت التطبيق وتحميل كل الملازم مجانا وبسرعه

MR

حمل التطبيق الأن

GET IT ON Google Play

ملازم ابتدائي | ملازم اعدادي | ملازم ثانـــوي | كورسات ونتائج امتحانات

موقع الدكتور محمد رزق
Dr.MoRizK